

JOURNAL OF COMBINATORIAL THEORY **11**, 169–180 (1971)On the 1-Factors of n -Connected Graphs*

JOSEPH ZAKS

*University of Washington, Seattle, Washington 98195,
and Wayne State University, Detroit, Michigan 48202**Communicated by F. Harary*

Received June 1969

L. W. Beineke and M. D. Plummer have recently proved [1] that every n -connected graph with a 1-factor has at least n different 1-factors. The main purpose of this paper is to prove that every n -connected graph with a 1-factor has at least as many as $n(n-2)(n-4) \dots 4 \cdot 2$, (or: $n(n-2)(n-4) \dots 5 \cdot 3$) 1-factors. The main lemma used is: if a 2-connected graph G has a 1-factor, then G contains a vertex V (and even two such vertices), such that each edge of G , incident to V , belongs to some 1-factor of G .

I. INTRODUCTION AND STATEMENT OF RESULTS. We consider here finite graphs, without loops or double edges. A 1-factor of a graph G is a subgraph of G , which contains all the vertices of G , each with valence 1.

Let $f(n)$ denote the maximum k for which the following is true: "If G is an n -connected graph with a 1-factor then G has at least k (different) 1-factors."

One can now reformulate Theorem 1 of [1] simply as $f(2) \geq 2$, and Theorem 2 of [1] becomes $f(n) \geq n$, for all $n \geq 2$.

Our extension of [1] is the following

THEOREM 2. $f(n) \geq n(n-2)(n-4) \dots 4 \cdot 2$ for even n , while for odd n , $f(n) = n(n-2)(n-4) \dots 5 \cdot 3$.

Let $F(G)$ denote the subgraph of G , which is the union of all the 1-factors of G . A vertex V of G will be called *totally covered by $F(G)$* if all the edges of G , incident to V , belong to $F(G)$.

The following is our main idea:

LEMMA 4. *If G is a 2-connected graph with a 1-factor, then G contains a vertex V which is totally covered by $F(G)$.*

* This paper was presented to the American Mathematical Society on April 26, 1969, and to the Calgary Combinatorial Conference in June, 1969.

To the proof of this lemma we have the following

COROLLARY 1. *If G is a 2-connected graph with a 1-factor, then G contains two vertices which are totally covered by $F(G)$.*

A simpler proof of Theorem 1 of [1] is given here, due to B. Grünbaum.

II. TERMINOLOGY. A graph G is *connected* if for each pair of its vertices V and W , G contains a $V - W$ path P : an ordered collection $(V_1, E_1, V_2, E_2, \dots, E_k, V_{k+1})$ of vertices V_i and edges E_i of G , such that $V_1 = V$, $V_{k+1} = W$ and for all $1 \leq i \leq k$, E_i is the edge (V_i, V_{i+1}) of G .

A graph G is *n -connected between two of its vertices V and W* if G contains n $V - W$ paths P_1, \dots, P_n such that $P_i \cap P_j = \{V, W\}$ for all $i \neq j$, $1 \leq i, j \leq n$. G is *n -connected* if it is n -connected between each pair of its vertices, and it contains at least two vertices.

It is well known that G is n -connected if it contains at least $n + 1$ vertices, and the deletion of no set of $n - 1$ or fewer vertices disconnects G .

Clearly, if G is n -connected, having a vertex V and an edge E , then both $G - V$ and $G - E$ are $n - 1$ connected.

The following is a well known property; the proof is added for completeness:

LEMMA 1. *If G is a 2-connected graph, and $E = (V_1, V_2)$ is an edge of G , such that $G - E$ is only 1-connected, then $G - E$ is only 1-connected between V_1 and V_2 .*

Proof. Let U and W be two vertices of G (with at most one of them being V_1 or V_2) such that G is 2-connected between U and W , and $G - E$ is only 1-connected between U and W . Let α_1 and α_2 be two disjoint (except for common end points) $U - W$ paths in G . Therefore E belongs to one of α_1 and α_2 , say to α_1 . $G - E$ is only 1-connected between U and W , therefore $G - E$ has a cut vertex X such that U and W belong to different connected components of $G - \{E, X\}$. Since α_2 connects U and W in $G - E$, $X \in \alpha_2$.

Since $E \in \alpha_1$, U is path-connected to one end point of E , in $G - \{E, X\}$, while W is path-connected to the other end point of E .

If $G - E$ is 2-connected between V_1 and V_2 , there are two disjoint (except for common end points) $V_1 - V_2$ -paths in $G - E$, say β_1 and β_2 .

X can be only on one of β_1 and β_2 , therefore V_1 and V_2 are connected in $G - \{E, X\}$, and therefore U and W are connected in $G - \{E, X\}$; this is a contradiction.

Therefore $G - E$ is not 2-connected between V_1 and V_2 , as promised by the lemma.

III. THEOREM 1 (Beineke-Plummer). *If a 2-connected graph G has a 1-factor, then G has at least another 1-factor.*

Proof (Grünbaum). The proof is by induction on the number of the edges, the assertion being obvious if G has 4 edges. Assuming the theorem is true for all the graphs with $< k$ edges, let G be a 2-connected graph with k edges and with a 1-factor F . If G has an edge, both end points of which are of valence 2, there is a trivial reduction, of both G and F .

Excluding this possibility, we observe that there necessarily exists an edge $E = (V_1, V_2)$ of $G - F$ both end points of which have valence ≥ 3 .

Let $G^* = G - E$. If G^* is 2-connected, then, since F is a 1-factor of G^* (as well as of G), it follows from the inductive assumption that G^* has at least two 1-factors, which are 1-factors of G , as promised. If G^* is not 2-connected, it has a cut vertex V ; $G^* - V$ has precisely two connected components, say G_1 and G_2 (with $V_i \in G_i$, $i = 1, 2$), since G is 2-connected.

If F contains the edge (V_1, V) , then the graph $G - G_2$, obtained by deleting G_2 from G , is 2-connected and has a 1-factor; hence it has, by the inductive assumption, at least one other 1-factor, which may be combined with $F \cap G_2$ to yield a new 1-factor of G . Similarly if (V_2, V) is an edge of F .

If neither of these cases arises, let us assume the notation is such that the edge of F which contains V has its other end point in G_1 . Let H_1 be the graph obtained from $G - G_2$ by adding the edge (V_1, V) (if it is not already there), and let H_2 be the graph obtained from $G - G_1$ by adding the edges (V_1, V_2) and (V_1, V) . Clearly, H_1 and H_2 are 2-connected, $F_1 = H_1 \cap F$ is a 1-factor of H_1 , while $F_2 = (V_1, V) \cup (H_2 \cap F)$ is a 1-factor of H_2 . Therefore there exist 1-factors F_i^* of H_i different from F_i , $i = 1, 2$.

If F_1^* does not contain the edge (V_1, V) then $(H_2 \cap F) \cup F_1^*$ is a 1-factor of G , different from F . If F_2^* does contain the edge (V_1, V) , then $[(F_2^* - (V_1, V)) \cup F_1]$ is a 1-factor of G different from F . If neither of these cases happens, then $(F_1^* \cup F_2^*) - (V_1, V)$ is the desired 1-factor of G .

This completes the proof of the theorem.

IV. With very little effort we can improve considerably the inequality $f(n) \geq n$, of [1], while using $f(2) \geq 2$, as follows:

LEMMA 2. *For each $n \geq 3$, $f(n) \geq f(n-1) + f(n-2)$.*

Proof. Let G be an n -connected graph with a 1-factor F , $n \geq 3$. Since G is 2-connected, it has by [1] another 1-factor, say F_1 . Let $E = (V_1, V_2)$ be an edge of G , such that $E \in F$ and $E \notin F_1$.

$G - E$ is $(n - 1)$ -connected, and has F_1 as a 1-factor, therefore $G - E$ has at least $f(n - 1)$ 1-factors, each one of which is a 1-factor of G , which does not contain E . $G^* = G - \{V_1, V_2\}$ is $(n - 2)$ -connected, and has $F - E$ as a 1-factor, hence G^* has at least $f(n - 2)$ 1-factors. Adding the edge E to a 1-factor of G^* yields a 1-factor of G ; therefore G has at least $f(n - 2)$ 1-factors, each one of which contains the edge E .

Clearly, these two collections of 1-factors of G are disjoint, therefore G has at least $f(n - 1) + f(n - 2)$ 1-factors.

This completes the proof of the lemma.

V. MAIN RESULTS. The main ideas of this paper are contained in the following two lemmas:

LEMMA 3. *If every n -connected graph with a 1-factor contains a vertex V , such that at least n edges of G , incident to V , belong to $F(G)$, then for each k , $k \geq 1$,*

$$f(k) \geq k(k - 2)(k - 4) \dots$$

Proof. The proof is by induction on k , as follows: for $k = 1$ this is the triviality that $f(1) \geq 1$; for $k = 2$ the claim is that $f(2) \geq 2$ which has been proved in [1] and in Section III here. To complete the proof it suffices to show that, for each k , $k \geq 3$,

$$f(k) \geq k \cdot f(k - 2).$$

Let G be a k -connected graph with a 1-factor. From the assumption of the lemma it follows that G contains vertices V, V_1, \dots, V_k and edges $E_i = (V, V_i)$, $1 \leq i \leq k$, such that each E_i belong to some 1-factor F_i of G .

Let $G_i = G - \{V, V_i\}$, for each $1 \leq i \leq k$. Then each graph G_i is $(k - 2)$ connected, and has a 1-factor: $F_i - E_i$. Let \tilde{F}_i be the collection of all the 1-factors of G_i ; therefore, $\text{card } \tilde{F}_i \geq f(k - 2)$. Let $F_i^* = \{A \cup E_i \mid A \in \tilde{F}_i\}$, for each $1 \leq i \leq k$. Clearly, each member of F_i^* is a 1-factor of G . Moreover, $F_i^* \cap F_j^* = \emptyset$ for $i \neq j$, $1 \leq i, j \leq k$, because each member of F_i^* contains the edge (V, V_i) and each member of F_j^* contains the edge (V, V_j) . Therefore G has at least $k \cdot f(k - 2)$ 1-factors, and the proof of Lemma 3 is completed.

In order to prove that the condition of Lemma 3 holds, we prefer to prove the following stronger claim:

LEMMA 4. *If G is a 2-connected graph with a 1-factor F , then G contains a vertex V which is totally covered by $F(G)$.*

Proof. The proof is by induction on the number k of the edges of G . The induction begins with $k = 4$, in which case $G = F(G)$, therefore all the vertices of G are totally covered by $F(G)$, because in this case G must be a (simple) cycle of length 4.

Let us assume that the assertion is true for all the appropriate graphs with less than k edges, and let G be a 2-connected graph with a 1-factor, having k edges.

To prove that G has a vertex, which is totally covered by $F(G)$, we suppose, on the contrary, that G has no such a vertex.

CLAIM 1. If G has an edge E , which belongs to one 1-factor F^* , but does not belong to another 1-factor F^{**} , of G , then $G - E$ is only 1-connected.

Proof. Suppose an edge E of G belongs to F^* , but not to F^{**} , and $G - E$ is 2-connected.

$G - E$ has $k - 1$ edges, and has a 1-factor F^{**} , hence, by the inductive assumption, $G - E$ contains a vertex V , which is totally covered by $F(G - E)$. Now, $E \in F^*$ therefore $E \in F(G)$, and clearly $F(G - E) \subset F(G)$; therefore the vertex V of G is totally covered by $F(G)$; this contradicts our supposition that G contains no such a vertex, and hence completes the proof of Claim 1.

Now, G is 2-connected and has a 1-factor F , therefore, by Theorem 1, G has another 1-factor F' . The subgraph $(F' - F) \cup (F - F')$ of G is not empty, and all of its vertices are of valence 2; therefore its connected components are (simple) cycles of length r , where r is even and $r \geq 4$.

Let $S = (V_1, E_1, V_2, E_2, \dots, V_r, E_r, V_1)$ be one such component, where $E_{2i} \in F$ and $E_{2i+1} \in F'$, $1 \leq 2i, 2i + 1 \leq r$.

For each $1 \leq i \leq r$, we have by Claim 1 that $G - E_i$ is only 1-connected, because each one of the edges E_i belongs to precisely one of the two 1-factors of G , F and F' .

Since no vertex of G is totally covered by $F(G)$, and clearly $S \subset F \cup F' \subset F(G)$, therefore each vertex V_i of S has valence ≥ 3 . Therefore there are edges E_1^*, \dots, E_r^* in G which are not edges of S , such that, for all $1 \leq i \leq r$, V_i is a vertex of E_i^* . Let $E_i^* = (V_i, V_i^*)$, for all $1 \leq i \leq r$.

G is 2-connected, therefore $G - V_i$, for each $1 \leq i \leq r$, is (at least) 1-connected. Let α_i be a path in $G - V_i$, connecting the vertex V_i^* with some vertex V_i^{**} of $S - V_i$, such that V_i^{**} is the only vertex of S on α_i ($1 \leq i \leq r$). Clearly $V_i^{**} \neq V_i$, and, if $V_i^* \in S$, then α_i is the degenerate path, consisting of the vertex V_i^* only, and then $V_i^{**} = V_i^*$.

CLAIM 2. V_i^{**} is not a neighbor, in S , of V_i , for all $1 \leq i \leq r$. (Remark that $r \geq 4$.)

Proof. Suppose, on the contrary, that, for some i , $1 \leq i \leq r$, V_i^{**} is a neighbor, in S , of V_i ; say $V_i^{**} = V_{i+1}$.[†] G is 2-connected, while $G - E_i$ is only 1-connected, hence by Lemma 1 $G - E_i$ is only 1-connected between V_i and V_{i+1} . However, $\alpha_i \cup E_i^*$ and $S - E_i$ are two disjoint $V_i - V_{i+1}$ paths in $G - E_i$. This is a contradiction, and therefore Claim 2 is true.

CLAIM 3. There is an index i for which $V_i V_{i+1} V_i^{**} V_{i+1}^{**}$ are in cyclic order on S .

Proof. Let i be an index for which the directed (in the sense of increasing i modulo r) arc from V_i to V_i^{**} on S contains the minimal number of intermediate vertices of S , among all such possible (directed) arcs. This minimal number is not zero, because V_i^{**} is different from V_i and from its two neighbors in S . If V_{i+1}^{**} were between V_{i+1} and V_i^{**} , the arc from V_{i+1} to V_{i+1}^{**} on S would contain less vertices than the arc from V_i to V_i^{**} , which contradicts the choice of i .

Therefore V_{i+1}^{**} is not between V_{i+1} and V_i^{**} , as required.

CLAIM 4. The index i , as given by Claim 3, is such that $G - E_i$ is 2-connected.

Proof. Suppose, on the contrary, that $G - E_i$ is not 2-connected. Since G is 2-connected, it follows, using Lemma 1, that $G - E_i$ is only 1-connected between V_i and V_{i+1} .

To prove that this is not the case, we will show that $G - E_i$ contains two $V_i - V_{i+1}$ paths, disjoint except for common end points, as follows:

If $\alpha_i \cap \alpha_{i+1} \neq \emptyset$, then $S - E_i$ is one such a path, while the other is contained in $\alpha_i \cup \alpha_{i+1} \cup E_i^* \cup E_{i+1}^*$.

If $\alpha_i \cap \alpha_{i+1} = \emptyset$, then $E_i^* \cup \alpha_i$ together with the part of S from V_{i+1} to V_i^{**} is one path, while the other one is $E_{i+1}^* \cup \alpha_{i+1}$, together with that part of S between V_{i+1}^{**} and V_i , which misses V_{i+1} . These two parts of S are disjoint because $V_i V_{i+1} V_i^{**} V_{i+1}^{**}$ are in cyclic order in S , as given by Claim 3.

The proof of Claim 4 is complete.

Clearly, Claims 1 and 4 are in contradiction, obtained by assuming that no vertex of G is totally covered by $F(G)$. The proof of Lemma 4 is complete.

Proof of Theorem 2. Theorem 2 states that, if an n -connected graph has a 1-factor, it has at least as many as $n(n-2)(n-4) \dots$ 1-factors, $n \geq 1$.

[†] $i+1$, as an index of a vertex of S , is $i+1 \pmod{r}$.

Since the assertion is trivial for $n = 1$, while it becomes just Theorem 1 for $n = 2$, we will assume that $n \geq 3$.

Every n -connected graph G with a 1-factor has, by Lemma 4, a vertex V which is totally covered by $F(G)$, since G is 2-connected. The valence of V is at least n , therefore the condition of Lemma 3 holds, and the proof of Theorem 2 follows immediately from Lemma 3.

To show that equality holds in $f(n) \geq n(n-2)(n-4) \dots 5 \cdot 3$, for odd n , observe that the complete graph with $n+1$ vertices, for odd n , has precisely $n(n-2)(n-4) \dots 5 \cdot 3$ 1-factors.

VI. REMARKS. We extend our Lemma 4 as follows:

COROLLARY 1. *If G is a 2-connected graph with a 1-factor, then G contains two vertices which are totally covered by $F(G)$.*

Proof. The proof is by induction on the number k of the edges of G , and is very similar to the proof of Lemma 4.

If $k = 4$, the only 2-connected graph with 1-factor having 4 edges is the cycle of length 4, which has all four of its vertices totally covered by its 1-factors.

Suppose our corollary is true for all the appropriate graphs with $k-1$ edges and let G be a 2-connected graph with k edges, such that G has a 1-factor F . G has by [1] another 1-factor, say F' .

Suppose now that G has at most one vertex, which is totally covered by $F(G)$.

CLAIM 1. *If G has an edge E , which belongs to one 1-factor F^* of G , but does not belong to another 1-factor F^{**} of G , then $G - E$ is only 1-connected.*

Proof. The proof is similar to the proof of Claim 1 in Lemma 4, except that here two vertices of G are under consideration of being totally covered by $F(G)$.

Next, let $S = (V_1, E_1, V_2, E_2, \dots, E_r, V_1)$ be a closed cycle, such that $E_{2i} \in F$ and $E_{2i+1} \in F'$, as in the proof of Lemma 4, $1 \leq 2i, 2i+1 \leq r$.

It follows from Claim 1 here that $G - E_i$ is 1-connected, for all $1 \leq i \leq r$.

If a vertex V_j of S is 2-valent in G , the two edges of G meeting at V_j are E_{j-1} and E_j , hence V_j is totally covered by $F(G)$. As a consequence of the assumption on G , at most one vertex of S is 2-valent (the rest have clearly valences ≥ 3).

If no vertex of S is 2-valent, the rest of the proof of Lemma 4 is applicable

(i.e., construct α_i , $1 \leq i \leq r$, and prove Claims 2, 3, and 4), thus contradicting the assumption on G .

In case one vertex of S , say V_1 , is 2-valent, we proceed as follows: since the valence of $V_i \geq 3$, for all $2 \leq i \leq r$, there are edges E_2^*, \dots, E_r^* of G which are not edges of S , such that V_i is a vertex of E_i^* , for all $2 \leq i \leq r$. Let $E_i^* = (V_i, V_i^*)$, $2 \leq i \leq r$. Since $G - V_i$ is connected, let α_i ($2 \leq i \leq r$) be a path in $G - V_i$, connecting V_i^* to some V_i^{**} on S , such that V_i^{**} is the only vertex of S on α_i ; if $V_i^* \in S$, then α_i degenerates to the vertex V_i^* and $V_i^{**} = V_1^*$. Clearly, $V_i^{**} \neq V_i$.

CLAIM 2. V_i^{**} is not a neighbor in S of V_i , for all $2 \leq i \leq r$.

The proof is as in the corresponding case in Lemma 4.

CLAIM 3. There is an index i , $2 \leq i \leq r - 1$, for which $V_i V_{i+1} V_i^{**} V_{i+1}^{**}$ are in cyclic order on S .

Proof. Observe that since V_1 is 2-valent, $V_i^{**} \neq V_1$ for all $2 \leq i \leq r$.

The directed arc, in the sense of increasing i modulo r , from V_2 to V_2^{**} on S does not contain V_1 . Let i , $2 \leq i \leq r$, be such that the directed arc from V_i to V_i^{**} on S does not contain V_1 and contains the minimal possible number of intermediate vertices.

If V_{i+1}^{**} belongs to the directed arc from V_i to V_i^{**} on S , then the directed arc from V_{i+1} to V_{i+1}^{**} on S does not contain V_1 and has less vertices than the directed arc from V_i to V_i^{**} on S ; this contradicts the choice of i .

Therefore V_{i+1}^{**} does not belong to the directed arc from V_i to V_i^{**} on S , and $V_i V_{i+1} V_i^{**} V_{i+1}^{**}$ are in cyclic order on S , as claimed.

CLAIM 4. The index i , as given by Claim 3, is such that $G - E_i$ is 2-connected.

The proof of this claim is the same as in the corresponding case in Lemma 4; Claims 1 and 4 contradict each other, hence the assumption on G is disproved, and G has two vertices which are totally covered by $F(G)$. The proof of Corollary 1 is complete.

If the valences of the vertices of a graph G are all $\geq k$ for $k \geq 2$ and V_1 and V_2 are vertices of G , then clearly the valences of the vertices of $G - \{V_1, V_2\}$ are all $\geq k - 2$. Therefore, using previous results implies

COROLLARY 2. If G is an n -connected graph, $n \geq 2$, G has a 1-factor and the valences of the vertices of G are all $\geq k$ (hence: $k \geq n$), then G has at least as many as the following number of 1-factors:

- (i) $k(k-2)(k-4) \dots (k-n+2)$ for even n , or
- (ii) $k(k-2)(k-4) \dots (k-n+3)$ for odd n .

Theorem 2 gives the precise value of $f(n)$, for all the odd n , while for the even n it gives only a lower bound.

Since every (simple) cycle of even length has precisely two 1-factors, it follows that $f(2) = 2$.

The party graph^{*} P_{2k} is defined as the complement of the graph consisting of k disjoint edges. P_6 is a 4-connected graph with precisely eight 1-factors, hence $f(4) \leq 8$, which together with Theorem 2 implies that $f(4) = 8$.

For even n , $n \geq 6$, we have the following:

$$\begin{aligned} 48 &\leq f(6) \leq 60, \\ 384 &\leq f(8) \leq 544, \\ 3840 &\leq f(10) \leq 6040, \text{ etc.,} \end{aligned}$$

where the upper bound for $f(n)$ is obtained using the graph P_{n+2} .

VII. CONJECTURES. Since $f(2k)$ was not determined here for $k \geq 3$, we would like to raise the following

CONJECTURE 1. $f(2k) > 2^k \cdot k!$, for all $k \geq 3$.

Relating to Corollary 1, we have

CONJECTURE 2. If G is an n -connected graph with a 1-factor, then G has n vertices which are totally covered by $F(G)$, for all $n \geq 3$.

In another direction we have

CONJECTURE 3. For every odd $n \geq 3$ and for $n = 4$, there is a unique graph G_n which is n -connected and has precisely $f(n)$ 1-factors.

Assuming the last conjecture is true, let $f^*(n)$ be the maximum k for which the following is true: "If G is an n -connected graph with a 1-factor, and $G \neq G_n$, then G has at least k (different) 1-factors."

CONJECTURE 4. $f^*(3) = 4, f^*(4) = 10$ and $f^*(5) = 30$, and $f^*(n) > f(n)$ for all odd $n \geq 3$ and $n = 4$.

^{*} P_{2k} "represents" a cocktail party of k couples, where everybody is talking to everybody else, except for each husband talking to his wife.

Let $f(n, v)$ be the maximum k for which the following is true: "If G is an n -connected graph with v vertices, and G has a 1-factor, then G has at least k (different) 1-factors." Let $B(n, k)$ denote the graph, obtained from the complete bipartite graph on n and $n + 2k$ vertices, by adding k disjoint edges with end points in the set of the $n + 2k$ vertices. $B(n, k)$ is n -connected and has $n!$ 1-factors, therefore $f(n, v) \leq n!$ for all $v \geq 2n$.

CONJECTURE 5. $f(n, v) = n!$, for all $v \geq 2n, n \geq 3$.

$B(n, 1)$ is a counter example to the following question:[†] "If G is an n -connected graph with a 1-factor, and n is large enough, does G have two 1-factors which have no edge in common?"

Most of the conjectures here are due to B. Grünbaum.

VIII. MAXIMAL MATCHINGS. A *matching* M of a graph G is a subgraph of G , which contains all the vertices of G , each with valence 0 or 1; M is called *maximal matching* if M is a matching and has the maximum possible number of edges. A vertex V of M with valence 0 in M is called *isolated* (in M).

A vertex V of a graph G is called *totally covered by* $M(G)$ if each edge of G , incident to V , belongs to a maximal matching of G .

Let $g(n)$ be the maximum k for which the following is true: "If G is an n -connected graph with no 1-factor, then G has at least k (different) maximal matchings."

We have the following

LEMMA 5. *If all the vertices of a graph G are of valence $\geq k$, and G has no 1-factors, then G has $k + 1$ vertices which are totally covered by $M(G)$.*

In particular, $n + 1$ vertices of an n -connected graph G with no 1-factors are totally covered by $M(G)$.

Lemma 5 implies

THEOREM 3. *For all $n \geq 3$, $g(n) \geq n \cdot g(n - 2) + \min\{g(n - 1), f(n - 1)\}$, and $g(1) = 2, g(2) = 3$.*

Proof of Lemma 5. Let G be a graph, all vertices of which have valence $\geq k$, and let M be a maximal matching of G . Since G has no 1-factors, M has an isolated vertex, call it V_0 . The valence of V_0 is $\geq k$, by assumption, so let V_1, \dots, V_t be all of the vertices of G such that (V_0, V_i) is an edge of G , for all $1 \leq i \leq t$, where $t \geq k$.

[†] Asked by W. T. Tutte, in private communication.

If V_i is an isolated vertex of M , for some $1 \leq i \leq t$, then $M \cup (V_0, V_i)$ is a matching of G , which has more edges than M has; this contradicts the maximality of M .

Therefore no vertex V_i , for $1 \leq i \leq t$, is isolated in M . Let us assume, without loss of generality, that the notation is such that (V_i, V_{i+1}) is an edge of M , for all odd i , $1 \leq i \leq 2p - 1$, for some integer p ; while (V_i, V_i^*) is an edge of M , for all $2p + 1 \leq i \leq t$, where V_i^* are vertices of G , such that $\{V_i \mid 1 \leq i \leq t\} \cap \{V_i^* \mid 2p + 1 \leq i \leq t\} = \emptyset$. Clearly, since the edges of a matching are disjoint, $V_i^* \neq V_j^*$ for all $i \neq j$ and $2p + 1 \leq i, j \leq t$.

We exhibit t additional matchings M_j of G , for all $1 \leq j \leq t$, as follows: for odd j and $1 \leq j \leq 2p - 1$, let $M_j = [M - (V_j, V_{j+1})] \cup (V_0, V_j)$; for even j and $2 \leq j \leq 2p$, let $M_j = [M - (V_{j-1}, V_j)] \cup (V_0, V_j)$; for all j , $2p + 1 \leq j \leq t$, let $M_j = [M - (V_j, V_j^*)] \cup (V_0, V_j)$.

Clearly, each M_j is a matching of G , and, since each one has the same number of edges as M has, M_j is a maximal matching of G , for all $1 \leq j \leq t$.

Since $(V_0, V_j) \in M_j$ for all $1 \leq j \leq t$, V_0 is totally covered by $M(G)$. We have established the following

LEMMA. *If V is an isolated vertex of a maximal matching M of the graph G under consideration, then V is totally covered by $M(G)$.*

Observing that V_{j+1} is an isolated vertex of M_j for all odd j and $1 \leq j \leq 2p - 1$; V_{j-1} is an isolated vertex of M_j for all even j and $2 \leq j \leq 2p$, and V_j^* is an isolated vertex of M_j for all $2p + 1 \leq j \leq t$, it follows from the last lemma that G has t more vertices which are totally covered by $M(G)$, and the proof of Lemma 5 is completed.

Proof of Theorem 3. It follows immediately from the proof of Lemma 5 that, if G is n -connected and has no 1-factors, then G has $n + 1$ different maximal matchings, hence $g(n) \geq n + 1$.

The graph consisting of two edges with a common vertex is 1-connected and has two maximal matchings, hence $g(1) = 2$; the cycle of length 3 is 2-connected and has three maximal matchings, hence $g(2) = 3$.

To prove the inequality in question, let $n \geq 3$ and let G be an n -connected graph with no 1-factors. It follows by Lemma 5 that G has a vertex V_0 totally covered by $M(G)$. Let (V_0, V_i) , for all $1 \leq i \leq t$, be all the edges of G , incident to V_0 . Since G is n -connected, $t \geq n$.

The graph $G - \{V_0, V_i\}$, for all $1 \leq i \leq t$, is $(n - 2)$ -connected and has no 1-factors, since adding the edge (V_0, V_i) to such a 1-factor yields a 1-factor of G , contradicting the assumption on G . However, if M^i is a maximal matching of $G - \{V_0, V_i\}$, $M^i \cup (V_0, V_i)$ is a maximal

matching of G , for all $1 \leq i \leq t$, and $M^i \neq M^j$ for $i \neq j$. Therefore G has at least $t \cdot g(n-2)$ maximal matchings, each one of which contains an edge of the form (V_0, V_i) , for a suitable i , $1 \leq i \leq t$.

In addition, G has maximal matchings having V_0 as an isolated vertex, as follows: $G - V_0$ is $(n-1)$ -connected and has either $f(n-1)$ 1-factors, or $g(n-1)$ maximal matchings (if it has no 1-factors). If \bar{M} is 1-factor of $G - V_0$, then $\bar{M} \cup \{V_0\}$ is a maximal matching of G of the required kind. If \bar{M} is a maximal matching of $G - V_0$ but $\bar{M} \cup V_0$ is not a maximal matching of G , then a maximal matching of G has one more edge than a maximal matching of $G - V_0$. However this contradicts the existence of a maximal matching of G having V_0 as an isolated vertex. Therefore $\bar{M} \cup V_0$ is a maximal matching of G of the required kind, hence G has at least $\min\{g(n-1), f(n-1)\}$ maximal matching with V_0 as an isolated vertex, and the proof of Theorem 3 is completed.

Except for the obvious questions that can be raised for the values of $g(n)$, $n \geq 3$, we state the following

CONJECTURE 6. For all $n \geq 3$, $g(n) > f(n)$.

ACKNOWLEDGMENTS

I would like to thank M. Rosenfeld for the valuable talks we have had, and, in particular, to express my great appreciation to Branko Grünbaum for his encouragement. The referee deserves my thanks for his many suggestions.

Added in proof: Conjecture 2 has been proved by L. Lovász (private communication).

REFERENCES

1. L. W. BEINEKE AND M. D. PLUMMER, On the 1-factors of a non-separable graph, *J. Combinatorial Theory* 2 (1967), 285-289.